

LIST OF PAPERS
authored or co-authored by Ernst Kenndler
29.11.2011

- [1]V.U. Weiss, X. Subirats, A. Pickl-Herk, G. Bilek, W. Winkler, M. Kumar, G. Allmaier, D. Blaas, and E. Kenndler, Characterization of Rhinovirus Subviral A-Particles via Capillary Electrophoresis, Electron Microscopy and Gas Phase Electrophoretic Mobility Molecular Analysis. Part I. Electrophoresis (submitted 24.11.2011).
- [2]X. Subirats, D. Blaas, and E. Kenndler, Recent developments of capillary and chip electrophoresis of biological particles: viruses, organelles, and cells. *Electrophoresis* 32 (2011) 1579–1590.
- [3]R. Mandrioli, E. Morganti, L. Mercolini, E. Kenndler, and M.A. Raggi, Fast analysis of amino acids in wine by capillary electrophoresis with laser-induced fluorescence detection. *Electrophoresis* 32 (2011) 2809-2815.
- [4]G. Bilek, N. Matschenko, A. Pickl-Herk, V.U. Weiss, X. Subirats, E. Kenndler, and D. Blaas, Liposomal Nanocontainers as Models for Viral Infection: Monitoring viral genomic RNA transfer through lipid membranes. *J. Virol.* 85 (2011) 8368-8375.
- [5]V.U. Weiss, G. Bilek, A. Pickl-Herk, X. Subirats, K. Niespodziana, R. Valenta, D. Blaas, and E. Kenndler, Liposomal leakage induced by virus-derived peptides, viral proteins and entire virions: rapid analysis by chip electrophoresis. *Anal. Chem.* 82 (2010) 8146-8152.
- [6]R. Mandrioli, A. Musenga, E. Kenndler, M. De Donno, M. Amore, and M.A. Raggi, Determination of Topiramate in human plasma by capillary electrophoresis with indirect UV detection. *J. Pharm. Biomed. Anal.* 53 (2010) 1319-1323.
- [7]E. Kenndler, Buffer type and concentration: mobility, selectivity, and resolution in CE. in: J. Cazes, (Ed.), *Encyclopedia of Chromatography*, CRC Press, Boca Raton, 2010, pp. 290-292.
- [8]E. Kenndler, CE in nonaqueous media. in: J. Cazes, (Ed.), *Encyclopedia of Chromatography*, CRC Press, Boca Raton, 2010, pp. 342-344.
- [9]E. Kenndler, Capillary isotachophoresis. in: J. Cazes, (Ed.), *Encyclopedia of Chromatography*, CRC Press, Boca Raton, 2010, pp. 298-299.
- [10]E. Kenndler, Organic solvents: influence on pKa. in: J. Cazes, (Ed.), *Encyclopedia of Chromatography*, CRC Press, Boca Raton, 2010, pp. 1691-1692.
- [11]E. Kenndler, Organic solvents: effect on ion mobility. in: J. Cazes, (Ed.), *Encyclopedia of Chromatography*, CRC Press, Boca Raton, 2010, pp. 1689-1690.
- [12]E. Kenndler, Organic solvents: classification for CE. in: J. Cazes, (Ed.), *Encyclopedia of Chromatography*, CRC Press, Boca Raton, 2010, pp. 1687-1688.
- [13]V.U. Weiss, G. Bilek, A. Pickl-Herk, D. Blaas, and E. Kenndler, Mimicking Virus Attachment to Host Cells employing Liposomes: Analysis by Chip Electrophoresis *Electrophoresis* 30 (2009) 2123-2128.
- [14]A. Téllez, and E. Kenndler, Formamide as organic modifier in micellar electrokinetic chromatography with SDS. *Electrophoresis* 30 (2009) 357-364.

- [15]A. Téllez, and E. Kenndler, Is the general conclusion justified that higher applicable field strength results in shorter analysis time with organic solvents in CE? *Electrophoresis* 30 (2009) 3978-3985.
- [16]M. Riesová, V. Hruška, E. Kenndler, and B. Gaš, Electromigration oscillations occur in ternary electrolyte systems with complex eigenmobilities as predicted by theory and detected by capillary electrophoresis. *J. Phys. Chem. B* 113 (2009) 12439-12446.
- [17]A. Musenga, M. Amore, R. Mandrioli, E. Kenndler, L. de Martino, and M.A. Raggi, Determination of duloxetine in human plasma by capillary electrophoresis with laser-induced fluorescence detection. *J. Chromatogr.B* 877 (2009) 1126-1132.
- [18]L. Kremser, D. Blaas, and E. Kenndler, Virus analysis using electromigration techniques. *Electrophoresis* 30 (2009) 133–140.
- [19]E. Kenndler, Organic solvents in CE. *Electrophoresis* 30 (2009) S101–S111.
- [20]G. Bilek, V.U. Weiss, A. Pickl-Herk, D. Blaas, and E. Kenndler, Chip electrophoretic characterization of liposomes with biological lipid composition: Coming closer to a model for viral infection. *Electrophoresis* 30 (2009) 4292-4299.
- [21]A.L. Valeri, L. Kremser, M.A. Raggi, B. Grüner, R. Vespalet, and E. Kenndler, Capillary electrophoresis of boron cluster compounds in aqueous and non-aqueous solvents. *Electrophoresis* 29 (2008) 1658–1666.
- [22]A. Téllez, V. Weiss, and E. Kenndler, An extended description of the effect of detergent monomers on migration in micellar electrokinetic chromatography. *Electrophoresis* 29 (2008) 3916 - 3923.
- [23]A. Schönemann, M. Eisbein, A. Unger, M. Dell'Mour, W. Frenzel, and E. Kenndler, Historic consolidants for wooden works of art in Saxony - an investigation by GC/MS and FTIR analysis. *Studies in Conservation* 53 (2008) 118-130.
- [24]A. Musenga, M.A. Saracino, D. Spinelli, E. Rizzato, G. Boncompagni, E. Kenndler, and M.A. Raggi, Analysis of the recent antipsychotic aripiprazole in human plasma by capillary electrophoresis and high performance liquid chromatography with diode array detection. *Analytica Chimica Acta* 612 (2008) 204–211.
- [25]A. Musenga, E. Kenndler, E. Morganti, F. Rasi, and M.A. Raggi, Analysis of the anti-Parkinson drug Pramipexole in human urine by capillary electrophoresis with laser-induced fluorescence detection. *Analytica Chimica Acta* 626 (2008) 89-96.
- [26]E. Kenndler, V.U. Weiss, G. Bilek, and L. Kremser, Chip-Elektrophorese von Viren. *GIT-Labor-Fachzeitschrift* 1 (2008) 16-18.
- [27]E. Kenndler, Analyse von Bindemitteln in Kunstobjekten mittels CE. *LCGC* (2008) 6-10.
- [28]A. Jouyban, and E. Kenndler, Impurity analysis of pharmaceuticals using capillary electromigration methods. *Electrophoresis* 29 (2008) 3531-3551.
- [29]C. Farina, L. Kremser, M.A. Raggi, and E. Kenndler, Determination of Ziprasidone in pharmaceutical formulations by capillary zone electrophoresis. *J. Pharm. Biomed. Anal.* 46 (2008) 471–476.
- [30]M. Dell'mour, A. Findeisen, I. Kaml, W. Baatz, and E. Kenndler, Capillary Electrophoresis as Analytical Method for the Characterisation of Natural Diterpenoid Resins in Artistic and Historic Works: A Comparison with Gas Chromatography. *The Open Analytical Chemistry Journal* 2 (2008) 67-73.
- [31]A.V. Colnaghi Simionato, E. Perin Moraes, E. Carrilho, M.F.M. Tavares, and E. Kenndler, Determination of amino acids by capillary electrophoresis-

- electrospray ionization-mass spectrometry: An evaluation of different protein hydrolysis procedures. *Electrophoresis* 29 (2008) 2051–2058.
- [32]V. Weiss, V. Kolivoška, L. Kremser, B. Gaš, D. Blaas, and E. Kenndler, Virus Analysis by Electrophoresis on a Microfluidic Chip. *J. Chromatogr.B* 860 (2007) 173–179.
- [33]J. Reijenga, L. Gagliardi, and E. Kenndler, Temperature dependence of acidity constant: a tool to affect separation selectivity in capillary electrophoresis. *J. Chromatogr.A* 1155 (2007) 142–145.
- [34]A. Musenga, R. Mandrioli, I. Comin, E. Kenndler, and M.A. Raggi, Determination of Vigabatrin in Human Plasma by Means of Capillary Electrophoresis with Laser-Induced Fluorescence Detection. *Electrophoresis* 28 (2007) 3535–3541.
- [35]A. Musenga, R. Mandrioli, P. Bonifazi, E. Kenndler, A. Pompei, and M.A. Raggi, Sensitive and selective determination of glutathione in probiotic bacteria by capillary electrophoresis-laser induced fluorescence. *Analyt. Bioanal. Chem.* 387 (2007) 917–924.
- [36]A. Musenga, E. Kenndler, L. Mercolini, M. Amore, S. Fanali, and M.A. Raggi, Determination of sertraline and N-desmethylsertraline in human plasma by capillary electrophoresis with laser-induced fluorescence detection. *Electrophoresis* 28 (2007) 1823–1831.
- [37]L. Mercolini, F. Bugamelli, E. Kenndler, G. Boncompagni, L. Franchini, and M.A. Raggi, Simultaneous determination of the antipsychotic drugs levomepromazine and clozapine and their metabolites in human plasma by a HPLC-UV method with solid phase extraction. *J. Chromatogr.B* 846 (2007) 273–280.
- [38]R. Martello, V. Kolivoška, M.A. Raggi, and E. Kenndler, Capillary electrophoresis of tricyclic antidepressant clomipramine and metabolites: electromigration and wall adsorption. *Electrophoresis* 28 (2007) 3650–3657.
- [39]L. Kremser, G. Bilek, and E. Kenndler, Effect of detergent on electromigration of proteins: capillary electrophoresis of VLDL receptor modules and viral proteins. *Electrophoresis* 28 (2007) 3684–3690.
- [40]L. Kremser, G. Bilek, D. Blaas, and E. Kenndler, Capillary electrophoresis of viruses, sub-viral particles and virus complexes. *J. Sep. Science* 30 (2007) 1704 – 1713.
- [41]V. Kolivoška, V. Weiss, L. Kremser, B. Gaš, D. Blaas, and E. Kenndler, Electrophoresis on a microfluidic chip for analysis of fluorescence labelled Human Rhinovirus. *Electrophoresis* 28 (2007) 4734–4740.
- [42]I. Kaml, and E. Kenndler, Characterization of natural organic binding media in museum objects by capillary electrophoresis. *Current Analytical Chemistry* 3 (2007) 33–40.
- [43]A. Findeisen, V. Kolivoška, I. Kaml, W. Baatsch, and E. Kenndler, Analysis of diterpenoic compounds in natural resins applied as binders in museum objects by capillary electrophoresis. *J. Chromatogr.A* 1157 (2007) 142–145.
- [44]F. Bugamelli, R. Mandrioli, E. Kenndler, C. Bartoletti, G. Boncompagni, and M.A. Raggi, Possible levomepromazine-clozapine interaction: two case reports. *Progress in Neuro-Psychopharmacology & Biological Psychiatry* 3 (2007) 567–570.
- [45]G. Bilek, L. Kremser, W. J., D. Blaas, and E. Kenndler, Mimicking Early Events of Virus Infection: Capillary Electrophoretic Analysis of Virus Attachment to Receptor - Decorated Liposomes. *Anal. Chem.* 79 (2007) 1620–1625.

- [46]A. Schönemann, W. Frenzel, A. Unger, and E. Kenndler, Investigation on the Fatty Acid Composition of Tung Oil for a Technical Examination of a Polychromed Wooden Altar. *Studies in Conservation* 51 (2006) 99-110.
- [47]M.A. Saracino, O. Gandolfi, R. Dall'Olio, L. Albers, E. Kenndler, and M.A. Raggi, Determination of Olanzapine in rat brain using liquid chromatography with coulometric detection and a rapid solid-phase extraction procedure. *J. Chromatogr. A* 1122 (2006) 21-27.
- [48]M. Mazanek, I. Kaml, and E. Kenndler, Capillary electrophoresis: an alternative to chromatography for analysis of natural organic binders. *Studies in Conservation* 51 (2006) 130-151.
- [49]R. Mandrioli, M.A. Saracino, S. Ferrari, D. Berardi, E. Kenndler, and M.A. Raggi, HPLC analysis of the second-generation antidepressant sertraline and its main metabolite N-desmethylsertraline in human plasma. *J. Chromatogr.B* 836 (2006) 116–119.
- [50]L. Kremser, M. Petsch, D. Blaas, and E. Kenndler, Influence of detergent additives for capillary electrophoresis of native and subviral rhinovirus particles. *Electrophoresis* 27 (2006) 1112-1121.
- [51]L. Kremser, M. Petsch, D. Blaas, and E. Kenndler, Capillary electrophoresis of affinity complexes between subviral 80S-particles of human rhinovirus and monoclonal antibody 2G2. *Electrophoresis* 27 (2006) 2630-2637.
- [52]E. Kenndler Capillary Electrophoresis. in: M. Akay, (Ed.), *Encyclopedia of Biomedical Engineering*, Wiley, New York, 2006, pp. 682-690.
- [53]A. Jouyban, and E. Kenndler, Theoretical and empirical approaches to express the mobility of small ions in capillary electrophoresis. *Electrophoresis* 27 (2006) 992–1005.
- [54]G. Bilek, L. Kremser, D. Blaas, and E. Kenndler, Analysis of liposomes by capillary electrophoresis and their use as carrier in electrokinetic chromatography. *J. Chromatogr.B* 841 (2006) 38-51.
- [55]G. Bilek, L. Kremser, D. Blaas, and E. Kenndler, Capillary electrophoresis of liposomes functionalised for protein binding. *Electrophoresis* 27 (2006) 3999–4007.
- [56]K. Vceláková, I. Zusková, S.P. Porras, B. Gaš, and E. Kenndler, Analyte and system eigenpeaks in nonaqueous capillary zone electrophoresis: Theoretical description and experimental confirmation with methanol as solvent. *Electrophoresis* 26 (2005) 463–472.
- [57]X. Subirats, S. Reinstadler, S.P. Porras, M.A. Raggi, and E. Kenndler, Comparison of methanol and acetonitrile as solvents for the separation of sertindole and its major metabolites by capillary zone electrophoresis. *Electrophoresis* 26 (2005) 3315–3324.
- [58]X. Subirats, S.P. Porras, M. Roses, and E. Kenndler, Nitromethane as solvent in capillary electrophoresis. *J. Chromatogr.A* 1079 (2005) 246-253.
- [59]C. Simó, M. Herrero, C. Neusüß, M. Pelzing, E. Kenndler, C. Barbas, E. Ibáñez, and A. Cifuentes, Characterization of proteins from *Spirulina platensis* microalga using capillary electrophoresis-ion trap-mass spectrometry and capillary electrophoresis-time of flight-mass spectrometry. *Electrophoresis* 26 (2005) 2674-2683.
- [60]S.P. Porras, and E. Kenndler, Are the asserted advantages of organic solvents in capillary electrophoresis real? A critical discussion. *Electrophoresis* 26 (2005) 3203-3220.
- [61]S.P. Porras, and E. Kenndler, Capillary electrophoresis in N,N-dimethylformamide. *Electrophoresis* 26 (2005) 3279–3291.

- [62]M. Petsch, B.X. Mayer-Helm, R. Sauermann, C. Joukhadar, and E. Kenndler, Determination of fosfomycin in pus by capillary zone electrophoresis. *J. Chromatogr.A* 1081 (2005) 55-59.
- [63]A. Nicodemou, M. Petsch, T. Konecsni, L. Kremser, E. Kenndler, J. Casasnovas, and D. Blaas, Rhinovirus-stabilizing activity of artificial VLDL-receptor variants defines a new mechanism for virus neutralization by soluble receptors. *FEBS Letters* 579 (2005) 5507–5511.
- [64]S.E. Lucangioli, V. Tripodi, E. Masrian, S.L. Scioscia, C.N. Carducci, and E. Kenndler, Enantioselective separation of rivastigmine by capillary electrophoresis with cyclodextrines. *J. Chromatogr.A* 1081 (2005) 31-35.
- [65]M. Herrero, D. Arráez-Román, A. Segura, E. Kenndler, B. Gius, M.A. Raggi, E. Ibáñez, and A. Cifuentes, Pressurized liquid extraction - capillary electrophoresis -mass spectrometry for the analysis of polar antioxidants in rosemary extracts. *J. Chromatogr.A* 1084 (2005) 54-62.
- [66]S. Harrison, I. Kaml, F. Rainer, and E. Kenndler, Identification of drying oils in mixtures of natural binding media used for artistic and historic works by capillary electrophoresis. *J. Sep. Sci.* 28 (2005) 1587–1594.
- [67]S. Harrison, I. Kaml, V. Prokoratova, M. Mazanek, and E. Kenndler, Animal glues in mixtures of natural binding media for artistic and historic objects: identification by capillary zone electrophoresis. *Anal. Bioanal. Chem.* 382 (2005) 1520–1526.
- [68]M. Größl, S. Harrison, I. Kaml, and E. Kenndler, Characterisation of natural polysaccharides (plant gums) used as binding media for artistic and historic works by capillary zone electrophoresis. *J. Chromatogr.A* 1077 (2005) 80-89.
- [69]K. Vceláková, I. Zusková, E. Kenndler, and B. Gaš, Determination of cationic mobilities and pKa values of twenty-two amino acids by capillary zone electrophoresis. *Electrophoresis* 25 (2004) 309-317.
- [70]I. Surowiec, I. Kaml, and E. Kenndler, Analysis of drying oils as binding media for objects of art by capillary zone electrophoresis with indirect UV and conductivity detection. *J. Chromatogr.A* 1024 (2004) 245-254.
- [71]V. Pucci, F. Bugamelli, R. Mandrioli, A. Ferranti, E. Kenndler, and M.A. Raggi, High-performance liquid chromatographic determination of Levetiracetam in human plasma: comparison of different sample clean-up procedures. *Biomed. Chromatogr.* 18 (2004) 37-44.
- [72]S.P. Porras, and E. Kenndler, Capillary zone electrophoresis in non-aqueous solutions: pH of the background electrolyte. *J. Chromatogr.A* 1037 (2004) 455-465.
- [73]S.P. Porras, and E. Kenndler, Formamide as solvent for capillary zone electrophoresis. *Electrophoresis* 25 (2004) 2946-2958.
- [74]M. Petsch, B.X. Mayer-Helm, R. Sauermann, C. Joukhadar, and E. Kenndler, Capillary electrophoresis analysis of fosfomycin in biological fluids for clinical pharmacokinetic studies. *Electrophoresis* 25 (2004) 2292-2298.
- [75]V.M. Okun, and E. Kenndler, Affinity Capillary Electrophoresis. in: A.S. Rathore, and A. Guttman, (Eds.), *Electrokinetic Phenomena. Principles and Applications in Analytical Chemistry and Microchip Technology*, Marcel Dekker, New York, 2004.
- [76]A. Musenga, A. Schedle, U. Demelbauer, L. Kremser, M.A. Raggi, and E. Kenndler, Separation of ethoxylated Bisphenol A dimethacrylates (Bis-EMA) in dental composite after derivatisation to ionisable amines by capillary zone electrophoresis. *J. Chromatogr. A* 1034 (2004) 221-226.

- [77]L. Kremser, M. Petsch, D. Blaas, and E. Kenndler, Labeling of Capsid Proteins and Genomic RNA of Human Rhinovirus with Two Different Fluorescent Dyes for Selective Detection by Capillary Electrophoresis. *Anal. Chem.* 76 (2004) 7360-7365.
- [78]L. Kremser, V.M. Okun, A. Nicodemou, D. Blaas, and E. Kenndler, Binding of Fluorescent Dye to Genomic RNA Inside Intact Human Rhinovirus after Viral Capsid Penetration Investigated by Capillary Electrophoresis. *Anal. Chem.* 76 (2004) 882-887.
- [79]L. Kremser, T. Konecsni, D. Blaas, and E. Kenndler, Fluorescence labeling of human rhinovirus capsid for analysis by capillary electrophoresis. *Anal. Chem.* 76 (2004) 4175-4181.
- [80]L. Kremser, D. Blaas, and E. Kenndler, Capillary electrophoresis of biological particles: viruses, bacteria, and eukaryotic cells. *Electrophoresis* 25 (2004) 2282-2291.
- [81]T. Konecsni, L. Kremser, L. Snyers, C. Rankl, F. Kilar, E. Kenndler, and D. Blaas, Twelve receptor molecules attach per viral particle of human rhinovirus serotype 2 via multiple modules. *FEBS Letters* 568 (2004) 99-104.
- [82]E. Kenndler, and A. Rizzi, *Electrokinetic Chromatography*. in: E. Heftmann, (Ed.), *Chromatography; Fundamentals and applications of chromatography and related differential migration methods; Part A: Fundamentals and techniques*, Elsevier, Amsterdam, 2004.
- [83]I. Kaml, K. Vcelakova, and E. Kenndler, Characterisation and identification of proteinaceous binding media (animal glues) from their amino acid profile by capillary zone electrophoresis. *J. Sep. Sci.* 27 (2004) 161-166.
- [84]B. Gaš, and E. Kenndler, System peaks in capillary zone electrophoresis. *Electrophoresis* 25 (2004) 3901-3912.
- [85]A. Schedle, M. Ivanova, and E. Kenndler, Determination of ethoxylated Bisphenol A dimethacrylate (BIS-EMA) monomers in dental composites by micellar electrokinetic chromatography. *J. Chromatogr.A* 990 (2003) 231-237.
- [86]O.L. Sánchez Muñoz, E. Pérez Hernández, M. Lämmerhofer, E. Tobler, W. Lindner, and E. Kenndler, Estimation and comparison of zeta potentials of silica-based anion exchange type porous particles for capillary electrochromatography from electrophoretic and electroosmotic mobility. *Electrophoresis* 24 (2003) 390-398.
- [87]V. Pucci, E. Kenndler, and M.A. Raggi, Quantitation of Oxcarbazepine and its Metabolites in Human Plasma by Micellar Electrokinetic Chromatography. *Biomed. Chromatogr.* 17 (2003) 231-238.
- [88]S.P. Porras, K. Sarmini, S. Fanali, and E. Kenndler, Medium effect (transfer activity coefficient) of methanol and acetonitrile on beta-cyclodextrin/benzoate complexation in capillary zone electrophoresis. *Anal. Chem.* 75 (2003) 1645-1651.
- [89]S.P. Porras, M.-L. Riekkola, and E. Kenndler, The principles of migration and dispersion in capillary zone electrophoresis in nonaqueous solvents. *Electrophoresis* 24 (2003) 1485-1498.
- [90]S.P. Porras, E. Marziali, B. Gaš, and E. Kenndler, Influence of Solvent on Temperature and Thermal Peak Broadening in Capillary Zone Electrophoresis. *Electrophoresis* 24 (2003) 1553-1564.
- [91]U.L. Peri-Okonny, E. Kenndler, R.J. Stubbs, and N.A. Guzman, Characterization of pharmaceutical drugs by a modified nonaqueous capillary electrophoresis-mass spectrometry method. *Electrophoresis* 24 (2003) 139-150.

- [92]I. Peric, and E. Kenndler, Recent developments in capillary electrokinetic chromatography with replaceable charged pseudostationary phases or additives. *Electrophoresis* 24 (2003) 2924-2934.
- [93]R. Mandrioli, N. Ghedini, F. Albani, E. Kenndler, and M.A. Raggi, Liquid chromatographic determination of oxcarbazepine and its metabolites in plasma of epileptic patients after solid-phase extraction. *J. Chromatogr.B* 783 (2003) 253-263.
- [94]S.E. Lucangioli, E. Kenndler, A. Carlucci, V.P. Tripodi, S.L. Scioscia, and C.N. Carducci, Relation between retention factors of immunosuppressive drugs in microemulsion electrokinetic chromatography with biosurfactants and octanol-water partition coefficients. *J. Pharm. Biomed. Anal.* 33 (2003) 871-878.
- [95]S.E. Lucangioli, C.N. Carducci, S.L. Scioscia, A. Carlucci, C. Bregni, and E. Kenndler, Comparison of the retention characteristics of different pseudostationary phases for microemulsion and micellar electrokinetic chromatography of betamethasone and derivatives. *Electrophoresis* 24 (2003) 984-991.
- [96]A. Jouyban, H.-K. Chan, B.J. Clark, and E. Kenndler, Comparison of different algorithms to calculate electrophoretic mobility of analytes as function of binary solvent composition. *Electrophoresis* 24 (2003) 1596-1602.
- [97]M. Ivanova, A. Piunti, E. Marziali, N. Komarova, M.A. Raggi, and E. Kenndler, Microemulsion electrokinetic chromatography applied for separation of levetiracetam from other antiepileptic drugs in polypharmacy. *Electrophoresis* 24 (2003) 992-998.
- [98]O.L. Sánchez Muñoz, O. Capote Peña, E. Dale Ledger, J.J. Llovera González, and E. Kenndler, Zeta potential of staphylococcal protein A (Z-domain) used for determination of complex constant with immunoglobulin Fc fragment by affinity electrophoresis in capillary format. *J. Cap. Elec.* 7 (2002) 31-36.
- [99]M.A. Raggi, V. Pucci, A. Maurizio, J. Muzikár, and E. Kenndler, Separation of carbamazepine and five metabolites, and analysis in human plasma by micellar electrokinetic capillary chromatography. *J. Chromatogr.B* 770 (2002) 217-225.
- [100]S.P. Porras, M.-L. Riekkola, and E. Kenndler, Capillary electrophoresis of anionic analytes in methanol: Effect of counter-ions on electrophoretic mobility. *Electrophoresis* 23 (2002) 367-374.
- [101]V.M. Okun, S. Nizet, D. Blaas, and E. Kenndler, Kinetics of thermal denaturation of human rhinoviruses in the presence of anti-viral capsid binders analyzed by capillary electrophoresis. *Electrophoresis* 23 (2002) 896-902.
- [102]J. Muzikár, T. van de Goor, and E. Kenndler, The principle cause of lower plate numbers in capillary zone electrophoresis with organic solvents. *Anal. Chem.* 74 (2002) 434-439.
- [103]J. Muzikár, T. van de Goor, B. Gaš, and E. Kenndler, Determination of Electroosmotic Flow Mobility with a Pressure-Mediated Dual-Ion Technique for Capillary Electrophoresis with Conductivity Detection using Organic Solvents. *J. Chromatogr.A* 960 (2002) 199-208.
- [104]J. Muzikár, T. van de Goor, B. Gaš, and E. Kenndler, Propylene carbonate as non-aqueous solvent for capillary electrophoresis: mobility and ionisation constants for aliphatic amines. *Anal. Chem.* 74 (2002) 428-433.
- [105]J. Muzikár, T. van de Goor, B. Gaš, and E. Kenndler, Electrophoretic mobilities of large organic ions in propylene carbonate, N,N-dimethylformamide, N,N-dimethylacetamide, acetonitrile and methanol: determination by capillary electrophoresis in non-aqueous solvents. *Electrophoresis* 23 (2002) 375-382.

- [106] J. Muzikár, G. Rozing, T. van de Goor, C. Eberwein, and E. Kenndler, Analysis of dimeric cyanine nucleic acid dyes by capillary zone electrophoresis in N,N-dimethylacetamide as non-aqueous organic solvent. *J. Chromatogr.A* 950 (2002) 249-255.
- [107] E. Marziali, M.A. Raggi, N. Komarova, and E. Kenndler, Octakis-6-sulfato-gamma-cyclodextrin as additive for capillary electrokinetic chromatography of dibenzoazepines: carbamazepine, oxcarbamazepine and their metabolites. *Electrophoresis* 23 (2002) 3020-3026.
- [108] A. Jouyban, S.C. Grosse, M.W. Coleman, H.K. Chan, E. Kenndler, and B.J. Clark, Calculation of electrophoretic mobility in mixed solvent buffers in capillary zone electrophoresis using a mixture response surface method. *Analyst* 127 (2002) 1188 - 1192.
- [109] M. Ivanova, E. Marziali, M.A. Raggi, and E. Kenndler, Microemulsion Electrokinetic Chromatography for the Separation of Carbamazepine, Oxcarbazepine and their Metabolites. *J. Sep. Science* 25 (2002) 863-871.
- [110] B. Gaš, and E. Kenndler, Peak broadening in microchip electrophoresis: A discussion of the theoretical background. *Electrophoresis* 23 (2002) 3817-3826.
- [111] F. Bugamelli, C. Sabbioni, R. Mandrioli, E. Kenndler, F. Albani, and M.A. Raggi, Simultaneous analysis of six antiepileptic drugs and two selected metabolites in human plasma by liquid chromatography after solid-phase extraction. *Anal. Chim. Acta* 472 (2002) 1-10.
- [112] U. Schneider, and E. Kenndler, Identification of plant and animal glues in museum objects by GC-MS, after catalytic hydrolysis of the proteins by the use of a cation exchanger, with simultaneous separation from the carbohydrates. *Fresenius J. Analyt. Chem.* 371 (2001) 81-87.
- [113] S.P. Porras, M.-L. Riekkola, and E. Kenndler, Capillary zone electrophoresis of basic drugs in non-aqueous acetonitrile with buffers based on a conventional pH scale. *Chromatographia* 53 (2001) 290-294.
- [114] S.P. Porras, M.-L. Riekkola, and E. Kenndler, Capillary zone electrophoresis of basic analytes in methanol as non-aqueous solvent: mobility and ionisation constant. *J. Chromatogr. A* 905 (2001) 259-268.
- [115] S.P. Porras, M.-L. Riekkola, and E. Kenndler, Electrophoretic mobilities of cationic analytes in non-aqueous methanol, acetonitrile and their mixtures: Influence of ionic strength and ion pair formation. *J. Chromatogr.A* 924 (2001) 31-42.
- [116] S.P. Porras, M.-L. Riekkola, and E. Kenndler, Resolution in capillary electrophoresis with nonaqueous methanol as solvent: Theoretical prediction and experimental confirmation. *Electrophoresis* 22 (2001) 3798-3804.
- [117] S.P. Porras, P. Jyske, M.-L. Riekkola, and E. Kenndler, Mobility and ionisation constant of basic drugs in methanol. Application of non-aqueous background electrolyte solutions for capillary zone electrophoresis based on a conventional pH scale. *J. Microcolumn Sep.* 13 (2001) 149-155.
- [118] V.M. Okun, R. Moser, B. Ronacher, E. Kenndler, and D. Blaas, VLDL Receptor Fragments of various lengths bind to human rhinovirus HRV2 with different stoichiometry: An Analysis of virus - receptor complexes by Capillary electrophoresis. *J. Biolog. Chem.* 276 (2001) 1057-1062.
- [119] V.M. Okun, R. Moser, D. Blaas, and E. Kenndler, Complexes between Monoclonal Antibodies and Receptor Fragments with a Common Cold Virus: Determination of Stoichiometry by Capillary Electrophoresis. *Anal. Chem.* 73 (2001) 3900-3906.

- [120] J. Muzikár, T. van de Goor, B. Gaš, and E. Kenndler, Extension of the Application Range of Organic Solvents in Capillary Electrophoresis by the Use of a Contactless Conductivity Detector. *J. Chromatogr.A* 924 (2001) 147-154.
- [121] S.E. Lucangioli, C.N. Carducci, V.P. Tripodi, and E. Kenndler, Retention of bile salts in micellar electrokinetic chromatography: relation of capacity factor to octanol-water partition coefficient and critical micellar concentration. *J. Chromatogr.B* 765 (2001) 113-120.
- [122] E. Kenndler, and D. Blaas, Capillary electrophoresis of macromolecular biological assemblies: bacteria and viruses. *TrAC* 20 (2001) 543-551.
- [123] E. Kenndler, Capillary Isotachophoresis. in: J. Cazes, (Ed.), *Encyclopedia of Chromatography*, M. Dekker, New York, 2001, pp. 126-127.
- [124] E. Kenndler, Capillary Electrophoresis in Nonaqueous Media. in: J. Cazes, (Ed.), *Encyclopedia of Chromatography*, M. Dekker, New York, 2001, pp. 117-120.
- [125] E. Kenndler, Classification of Organic Solvents in Capillary Electrophoresis. in: J. Cazes, (Ed.), *Encyclopedia of Chromatography*, M. Dekker, New York, 2001, pp. 178-179.
- [126] E. Kenndler, Effect of Organic Solvents on Ion Mobility. in: J. Cazes, (Ed.), *Encyclopedia of Chromatography*, M. Dekker, New York, 2001, pp. 272-274.
- [127] E. Kenndler, Influence of Organic Solvents on *pKa*. in: J. Cazes, (Ed.), *Encyclopedia of Chromatography*, M. Dekker, New York, 2001, pp. 433-435.
- [128] E. Kenndler, Buffer Type and Concentration, Effect on Mobility, Selectivity, and Resolution in Capillary Electrophoresis. in: J. Cazes, (Ed.), *Encyclopedia of Chromatography*, M. Dekker, New York, 2001, pp. 98-100.
- [129] A. Jouybana, H.K. Chan, B.J. Clark, and E. Kenndler, Mathematical representation of electrophoretic mobility in mixed aqueous-methanolic buffers in capillary zone electrophoresis. *J. Microcolumn Sep.* 13 (2001) 345-350.
- [130] G. Izzo, A. Raggi, B. Maichel, and E. Kenndler, Separation of olanzapine, carbamazepine, and their main metabolites by capillary electrophoresis with pseudo-stationary phases. *J. Chromatogr.B* 752 (2001) 47-53.
- [131] K. Gogova, B. Maichel, E. Kenndler, and B. Gaš, Electrokinetic chromatography with micelles, polymeric and monomeric additives with similar chemical functionality as pseudo-stationary phases. *J. Chromatogr.A* 916 (2001) 79-87.
- [132] M.A. Raggi, G. Casamenti, R. Mandrioli, G. Izzo, and E. Kenndler, Quantitation of olanzapine in tablets by high performance liquid chromatography, capillary zone electrophoresis, derivative spectrometry or linear voltammetry. *J. Pharm. Biomed. Anal.* 23 (2000) 973-981.
- [133] E. Pucci, M. Raggi, and E. Kenndler, Quality Control of Pharmaceutical Formulations of Neuroleptic Drugs by Capillary Zone Electrophoresis. *J. Liq. Chrom.* 23 (2000) 25-34.
- [134] B. Potocek, E. Chmela, B. Maichel, E. Tesarova, E. Kenndler, and B. Gaš, Capillary Electrokinetic Chromatography with Charged Linear Polymers as a Nonmicellar PseudoStationary Phase: Determination of Capacity Factors and Characterization by Solvation Parameters. *Anal. Chem.* 72 (2000) 74-80.
- [135] V.M. Okun, B. Ronacher, D. Blaas, and E. Kenndler, Capillary Zone Electrophoresis with Post-Column Infectivity Assay for the Analysis of Different Serotypes of Human Rhinovirus (Common Cold Virus). *Anal. Chem.* 72 (2000) 2553-2558.
- [136] V.M. Okun, B. Ronacher, D. Blaas, and E. Kenndler, Affinity capillary electrophoresis for the assessment of complex formation between viruses and monoclonal antibodies. *Anal. Chem.* 72 (2000) 4634-4639.

- [137]B. Maichel, and E. Kenndler, Recent innovations in capillary electrokinetic chromatography with replaceable charged pseudostationary phases or additives. *Electrophoresis* 21 (2000) 3160-3173.
- [138]B. Maichel, K. Gogova, B. Gaš, and E. Kenndler, Comparison of separation selectivity in capillary electrokinetic chromatography using cationic linear polymeric pseudo-stationary phase or monomeric additives of similar structure. *J. Chromatogr.A* 894 (2000) 25-34.
- [139]B. Maichel, B. Gaš, and E. Kenndler, Diffusion coefficient and capacity factor in capillary electrokinetic chromatography with replaceable charged polymeric pseudophase. *Electrophoresis* 21 (2000) 1505-1512.
- [140]B. Gaš, and E. Kenndler, Dispersive phenomena in electromigration methods. *Electrophoresis* 21 (2000) 3888-3897.
- [141]T. Shimizu, and E. Kenndler, Capillary electrophoresis of small solutes in linear polymer solutions: relation between ionic mobility, diffusion coefficient and viscosity. *Electrophoresis* 20 (1999) 3364-3372.
- [142]K. Sarmini, G. Reich, and E. Kenndler, Similarity of binary mixed aqueous-organic solvents for capillary zone electrophoresis based on the ionic mobility of substituted benzoates evaluated by cluster analysis and information theory. *J. Microcolumn Sep.* 11 (1999) 576-581.
- [143]K. Sarmini, and E. Kenndler, Ionization constants of weak acids and bases in organic solvents. *J. Biochem. Biophys. Meth.* 38 (1999) 123-137.
- [144]K. Sarmini, and E. Kenndler, Capillary zone electrophoresis in mixed aqueous-organic media: Effect of organic solvent on actual ionic mobilities, acidity constants and separation selectivity of substituted aromatic acids. IV. Acetonitrile. *J. Chromatogr.A* 833 (1999) 245-259.
- [145]V. Pucci, M. Raggi, and E. Kenndler, Separation of eleven central nervous system drugs by capillary zone electrophoresis. *J. Chromatogr.B* 728 (1999) 263-271.
- [146]E. Pucci, M. Raggi, and E. Kenndler, Separation of antipsychotic drugs (clozapine, loxapine) and their metabolites by capillary zone electrophoresis. *J. Chromatogr.A* 853 (1999) 461-468.
- [147]V.M. Okun, B. Ronacher, D. Blaas, and E. Kenndler, Analysis of common cold virus (human rhinovirus serotype 2) by capillary zone electrophoresis: the problem of peak identification. *Anal. Chem.* 71 (1999) 2028-2032.
- [148]V.M. Okun, D. Blaas, and E. Kenndler, Separation and biospecific identification of subviral particles of human rhinovirus serotype 2 (HRV2) by capillary zone electrophoresis. *Anal. Chem.* 71 (1999) 4480-4485.
- [149]B. Maichel, B. Potoczek, B. Gaš, and E. Kenndler, Capillary electrokinetic chromatography with polyethyleneimine as replaceable cationic pseudo-stationary phase: influence of methanol and acetonitrile on separation selectivity. *J. Chromatogr.A* 853 (1999) 121-129.
- [150]K. Sarmini, and E. Kenndler, Capillary zone electrophoresis in mixed aqueous-organic media: Effect of organic solvent on actual ionic mobilities, acidity constants and separation selectivity of substituted aromatic acids. I. Methanol. *J. Chromatogr.A* 806 (1998) 325-335.
- [151]K. Sarmini, and E. Kenndler, Capillary zone electrophoresis in mixed aqueous-organic media: Effect of organic solvent on actual ionic mobilities, acidity constants and separation selectivity of substituted aromatic acids. II. Ethanol. *J. Chromatogr.A* 811 (1998) 201-209.
- [152]K. Sarmini, and E. Kenndler, Capillary zone electrophoresis in mixed aqueous-organic media: Effect of organic solvent on actual ionic mobilities, acidity

- constants and separation selectivity of substituted aromatic acids. III. Propanol. *J. Chromatogr.A* 818 (1998) 209-215.
- [153]K. Sarmini, and E. Kenndler, Capillary electrophoresis in mixed aqueous-organic solvents: Effect of aprotic dipolar N,N-dimethylformamide and N,N-dimethylacetamide on acidity constants of substituted aromatic acids. *J. Cap. Elec.* 5 (1998) 103-110.
- [154]B. Potocek, B. Maichel, B. Gaš, M. Chiari, and E. Kenndler, Separation of neutral compounds by capillary electrokinetic chromatography with a replaceable charged linear polymer as pseudo-stationary phase. *J. Chromatogr.A* 798 (1998) 269-273.
- [155]B. Maichel, B. Potocek, B. Gas, M. Chiari, and E. Kenndler, Separation of neutral compounds by capillary electrokinetic chromatography using polyethyleneimine as cationic pseudo-stationary phase. *Electrophoresis* 19 (1998) 2124-2128.
- [156]E. Kenndler, Theory of Capillary Zone Electrophoresis. in: M.G. Khaledi, (Ed.), *High Performance Capillary Electrophoresis, Theory, Techniques and Applications*, John Wiley, New York, 1998, pp. 25-76.
- [157]E. Kenndler, Dependence of Analyte Separation on Electroosmotic Flow in Capillary Zone Electrophoresis: Quantitative Description by the Reduced Mobility. *J. Microcolumn Sep.* 10 (1998) 273-279.
- [158]U. Schnabel, C.-H. Fischer, and E. Kenndler, Characterization of colloidal gold nano-particles according to size by capillary zone electrophoresis. *J. Microcol. Sep.* 9 (1997) 529-534.
- [159]K. Sarmini, and E. Kenndler, Influence of organic solvents on the separation selectivity of capillary electrophoresis. *J. Chromatogr. A* 792 (1997) 3-11.
- [160]M. Morand, D. Blaas, and E. Kenndler, Reduction of wall adsorption in capillary zone electrophoresis of a basic single chain antibody fragment by a cationic polymeric buffer additive. *J. Chromatogr.B* 691 (1997) 192-198.
- [161]M. Minarik, B. Gaš, and E. Kenndler, Size-based separation of polyelectrolytes by capillary zone electrophoresis with linear polymeric additives: Migration regimes and selectivity of poly(styrenesulphonates) in solutions of derivatized cellulose. *Electrophoresis* 18 (1997) 98-103.
- [162]B. Ingelse, K. Sarmini, J.C. Reijenga, E. Kenndler, and F. Everaerts, Chiral interactions in capillary zone electrophoresis: Computer simulation and comparison with experiment. *Electrophoresis* 18 (1997) 938-942.
- [163]B. Gaš, M. Stedry, and E. Kenndler, Peak broadening in capillary zone electrophoresis. *Electrophoresis* 18 (1997) 2123-2133.
- [164]C.-H. Fischer, and E. Kenndler, Analysis of colloids IX. Investigation of the electrical double layer of colloidal inorganic nm-particles by size exclusion chromatography. *J. Chromatogr.A* 773 (1997) 179-187.
- [165]P. Blatny, F. Kvasnicka, and E. Kenndler, Trace determination of iron in water on the ppb level by on-line coupling of capillary isotachophoresis and capillary zone electrophoresis with UV detection of the Fe(III)-EDTA complex. *J. Chromatogr.A* 757 (1997) 297-302.
- [166]M. Štedry, S. Popelka, B. Gaš, and E. Kenndler, Influence of transference numbers on migration and deformation of concentration boundaries in capillary zone electrophoresis. *Electrophoresis* 17 (1996) 1121-1125.
- [167]W. Schützner, S. Fanali, A. Rizzi, and E. Kenndler, Separation of diastereomers by capillary zone electrophoresis in free solution with polymer additive and organic solvent component : Effect of pH and solvent composition. *J. Chromatogr.A* 719 (1996) 411-420.

- [168]U. Schnabel, F. Groiss, D. Blaas, and E. Kenndler, Determination of the pI of Human Rhinovirus Serotype 2 by Capillary Isoelectric Focusing. *Anal. Chem.* 68 (1996) 4300-4303.
- [169]M. Minarik, F. Groiss, B. Gaš, D. Blaas, and E. Kenndler, Dispersion Effects accompanying Pressurized Zone Mobilization in Capillary Isoelectric Focusing of Proteins. *J. Chromatogr.A* 738 (1996) 123-128.
- [170]K. Kenndler-Blachkoltm, S. Popelka, B. Gas, and E. Kenndler, Apparent Baseline Irregularities for Neutral Markers in Capillary Zone Electrophoresis with Electroosmotic Flow. *J. Chromatogr. A* 734 (1996) 351-356.
- [171]E. Kenndler, Effect of electroosmotic flow on selectivity, efficiency, and resolution in capillary zone electrophoresis expressed by the dimensionless reduced mobility. *J. Cap. Elec.* 3 (1996) 191-198.
- [172]P. Blatny, F. Kvasnicka, and E. Kenndler, Determination of anionic feed additives by two dimensional capillary isotachophoresis. *J. Chromatogr.A* 737 (1996) 255-262.
- [173]M. Štedrý, B. Gaš, and E. Kenndler, Dynamics of peak dispersion in capillary zone electrophoresis including wall adsorption. II. Exact analysis of unsteady linear adsorptive dispersion. *Electrophoresis* 16 (1995) 2027-2033.
- [174]W. Schützner, S. Fanali, A. Rizzi, and E. Kenndler, Separation of diastereomers by capillary zone electrophoresis with polymer additives: effect of polymer type and chain length. *Anal. Chem.* 67 (1995) 3866-3870.
- [175]B. Potocek, B. Gaš, E. Kenndler, and M. Stedry, Electroosmosis in capillary zone electrophoresis with non-uniform zeta potential. *J. Chromatogr. A* 709 (1995) 51-62.
- [176]M. Minarik, B. Gaš, A. Rizzi, and E. Kenndler, Plate height contribution of wall adsorption in capillary electrophoresis of proteins. *J. Cap. Elec.* 2 (1995) 89-96.
- [177]B. Gaš, M. Stedry, A. Rizzi, and E. Kenndler, Dynamics of peak dispersion in capillary zone electrophoresis including wall adsorption. I. Theoretical Model and Results of Simulation. *Electrophoresis* 16 (1995) 958-967.
- [178]B. Gaš, M. Stedry, and E. Kenndler, Contribution of the electroosmotic flow to peak broadening in capillary zone electrophoresis with uniform zeta potential. *J. Chromatogr.A* 709 (1995) 63-68.
- [179]W. Friedl, J.C. Reijenga, and E. Kenndler, Ionic strength and charge number correction for mobilities of multivalent organic anions in capillary electrophoresis. *J. Chromatogr.A* 709 (1995) 163-170.
- [180]S. Cladrowa-Runge, R. Hirz, E. Kenndler, and A. Rizzi, Enantiomeric separation of amphetamine related drugs by capillary zone electrophoresis using native and derivatized α -cyclodextrin as chiral additives. *J. Chromatogr.A* 710 (1995) 339-345.
- [181]M. Chiari, and E. Kenndler, Capillary zone electrophoresis in organic solvents: separation of anions in methanolic buffer solutions. *J. Chromatogr.A* 716 (1995) 303-309.
- [182]P. Blatny, F. Kvasnicka, and E. Kenndler, Determination of Phytic Acid in Cereal Grains, Legumes and Feeds by Capillary Isotachophoresis. *J. Agr. Food Chem.* 43 (1995) 129-133.
- [183]P. Blatny, C.-H.- Fischer, A. Rizzi, and E. Kenndler, Linear polymers applied as pseudophases in capillary zone electrophoresis of azo-compounds used as textile dyes. *J. Chromatogr.A* 717 (1995) 157-166.
- [184]P. Blatny, C.-H. Fischer, and E. Kenndler, Improvement of the separation selectivity in capillary zone electrophoresis of synthetic cationic dyes by

- polyvinylpyrrolidon as pseudo-stationary phase. *Fresenius J. Anal. Chem.* 352 (1995) 712-714.
- [185] W. Schützner, G. Caponecchi, S. Fanali, A. Rizzi, and E. Kenndler, Improved separation of diastereomeric derivatives of enantiomers by a physical network of linear polyvinylpyrrolidone applied as pseudophase in capillary zone electrophoresis. *Electrophoresis* 15 (1994) 769-773.
- [186] J.C. Reijenga, and E. Kenndler, Computational Simulation of Migration and Dispersion in Free Capillary Zone Electrophoresis. II. Results of Simulation and Comparison with Measurements. *J. Chromatogr.A* 659 (1994) 417-426.
- [187] J.C. Reijenga, and E. Kenndler, Computational Simulation of Migration and Dispersion in Free Capillary Zone Electrophoresis. I. Description of the Theoretical Model. *J. Chromatogr.A* 659 (1994) 403-415.
- [188] E. Kenndler, and H. Poppe, Entangled, noncross-linked polymer solutions used as physical network in capillary zone electrophoresis. *J. Cap. Elec.* 1 (1994) 144-155.
- [189] B. Gassner, W. Friedl, and E. Kenndler, Wall adsorption of small anions in capillary zone electrophoresis induced by cationic trace constituents of the buffer. *J. Chromatogr.A* 680 (1994) 25-31.
- [190] W. Friedl, and E. Kenndler, Limitations of the optimization of the resolution by the buffer pH in capillary zone electrophoresis. *Fresenius J. Anal. Chem.* 348 (1994) 576-582.
- [191] P. Blatny, F. Kvasnicka, and E. Kenndler, Time course of formation of inositol phosphates during enzymatic hydrolysis of phytic acid (myo-inositol hexaphosphoric acid) by phytase determined by capillary isotachophoresis. *J. Chromatogr. A* 679 (1994) 345-348.
- [192] C. Schwer, B. Gaš, Lottspeich, and E. Kenndler, Computer Simulation and Experimental Evaluation of On-Column Sample Preconcentration in Capillary Zone Electrophoresis by Discontinuous Buffer Systems. *Anal. Chem.* 65 (1993) 2108-2115.
- [193] W. Schützner, S. Fanali, A. Rizzi, and E. Kenndler, Separation of diastereomeric derivatives of enantiomers by capillary zone electrophoresis with a polymer network: use of polyvinylpyrrolidone as buffer additive. *J. Chromatogr.* 639 (1993) 375-378.
- [194] E. Kenndler, Organic Solvents in Capillary Electrophoresis. in: N.A. Guzman, (Ed.), *Capillary Electrophoresis Technology*, Marcel Dekker, New York, Basel, Hongkong, 1993, pp. 161-186.
- [195] J.F.K. Huber, E. Kenndler, G. Reich, W. Hack, and J. Wolf, Optimal Selection of Gas Chromatographic Columns for the Analytical Control of Chemical Warfare Agents by Application of Information Theory to Retention Data. *Anal. Chem.* 65 (1993) 2903-2906.
- [196] K. Futschik, M. Ammann, S. Bachmayer, and E. Kenndler, Determination of Ionic Species formed during Growth of *Escherichia coli* by Capillary Isotachophoresis. *J. Chromatogr.* 644 (1993) 389-395.
- [197] W. Friedl, and E. Kenndler, Resolution as a Function of the pH of the Buffer Based on the Analyte Charge Number for Multivalent Ions in Capillary Zone Electrophoresis without Electroosmotic Flow: Theoretical Prediction and Experimental Evaluation. *Anal. Chem.* 65 (1993) 2003-2009.
- [198] C. Schwer, and E. Kenndler, Peak Broadening in Capillary Zone Electrophoresis with Electro-Osmotic Flow: Dependence of Plate Number and Resolution on Charge Number. *Chromatographia* 33 (1992) 331-335.

- [199]W. Schützner, and E. Kenndler, Electrophoresis in Synthetic Organic Polymer Capillaries: Variation of Electroosmotic Velocity and Zeta Potential with pH and Solvent Composition. *Anal. Chem.* 64 (1992) 1991-1995.
- [200]E. Kenndler, and C. Schwer, Peak dispersion and separation efficiency in high-performance zone electrophoresis with gel-filled capillaries. *J.Chromatogr.* 595 (1992) 313-318.
- [201]E. Kenndler, K. Schmidt-BeiwI, F. Mairinger, and M. Pöhm, Identification of proteinaceous binding media of easel paintings by gas chromatography of the amino acid derivatives after catalytic hydrolysis by a protonated cation exchanger. *Fresenius J.Anal.Chem.* 342 (1992) 135-141.
- [202]E. Kenndler, and W. Friedl, Adjustment of resolution and analysis time in capillary zone electrophoresis by varying the pH of the buffer. *J.Chromatogr.* 608 (1992) 161-170.
- [203]A. Widhalm, C. Schwer, D. Blaas, and E. Kenndler, Capillary zone electrophoresis with a linear, non-cross-linked polyacrylamide gel: separation of proteins according to molecular mass. *J.Chromatogr.* 549 (1991) 446-451.
- [204]A. Widhalm, and E. Kenndler, Determination of morpholine and N-methylmorpholine formed as degradation products in cellulose fibre production by capillary isotachophoresis. *Anal.Chem.* 63 (1991) 645-648.
- [205]C. Schwer, and E. Kenndler, Electrophoresis in Fused-Silica Capillaries: The Influence of Organic Solvent on the Electroosmotic Velocity and the zeta Potential. *Anal.Chem.* 63 (1991) 1801-1807.
- [206]E. Kenndler, and C. Schwer, Nondependence of Diffusion-Controlled Peak Dispersion on Diffusion Coefficient and Ionic Mobility in Capillary Zone Electrophoresis without Electroosmotic Flow. *Anal.Chem.* 63 (1991) 2499-2502.
- [207]E. Kenndler, and K. Schmid-BeiwI, Effect of sodium dodecylsulphate in protein samples on the separation with free capillary zone electrophoresis. *J.Chromatogr.* 545 (1991) 397-402.
- [208]C. Schwer, and E. Kenndler, Capillary Electrophoresis. *Chromatographia* 30 (1990) 546-554.
- [209]E. Kenndler, C. Schwer, and D. Kaniansky, Purity Control of Riboflavin-5'-Phosphate (Vitamin B2-Phosphate) by Capillary Zone Electrophoresis. *J.Chromatogr.* 508 (1990) 203-207.
- [210]E. Kenndler, C. Schwer, and J.F.K. Huber, Determination of O,O-bis (2-ethylhexyl) phosphorodithioic acid in water at the ppt-level by GC/ECD after reactive extraction and pre-column derivatization. *Intern.J.Environ.Analyt.Chem.* 41 (1990) 1-13.
- [211]E. Kenndler, C. Schwer, B. Fritzsche, and M. Pöhm, Determination of arbutine in *uvae-ursi* folium (bearberry leaves) by capillary zone electrophoresis. *J.Chromatogr.* 514 (1990) 383-388.
- [212]E. Kenndler, and C. Schwer, Kapillarzonenelektrophorese. *GIT Fachz.Lab.* 10 (1990) 1241-1252.
- [213]E. Kenndler, and F. Mairinger, Examination of the diterpenoic resin components of anatomical wax models from the eighteenth century by gas chromatography. *Fresenius J.Anal.Chem.* 338 (1990) 635-640.
- [214]E. Kenndler, and B. Gassner, Cluster Analysis Applied to the Selection and Combination of Buffering Electrolyte Systems Used for Capillary Electrophoresis of Anions with Water or Methanol as Solvents. *Anal.Chem.* 62 (1990) 431-436.

- [215]E. Kenndler, Capillary Zone Electrophoresis and Isotachophoresis: A Comparison by Information Theory. *Chromatographia* 30 (1990) 713-718.
- [216]E. Kenndler, C. Schwer, and P. Jenner, Isotachophoresis in Mixed Solvents Consisting of Water, Methanol and Dimethylsulphoxide. III. Influence of the Solvent Composition on the Dissociation Constants and Mobilities of Non- and Hydroxy-substituted Aliphatic Carboxylic Acids. *J.Chromatogr.* 470 (1989) 57-68.
- [217]E. Kenndler, C. Schwer, and J.F.K. Huber, Determination of 1,2,4-Trimethylbenzene (Pseudocumene) in Serum of a Person Exposed to Liquid Scintillation Counting Solutions by GC/MS. *J.Anal.Toxicology* 13 (1989) 211-213.
- [218]E. Kenndler, and J.F.K. Huber, Methodische Fortschritte der Gaschromatographie. in: W.F. R.Borsdorf, K.Gürtler, (Ed.), *Analytiker Taschenbuch*, Springer Verlag, Berlin, Heidelberg, New York (London, Paris, Tokio),, 1989, pp. 3-33.
- [219]E. Kenndler, and G. Reich, Characterization and Selection of Electrolyte Systems for Isotachophoresis of Anions by Cluster Analysis. *Anal.Chem.* 60 (1988) 120-124.
- [220]E. Kenndler, F. Mairinger, W.P. Bauer, F. Rainer, and K. Schmidt-BeiwI, Technical Examination and Restoration of a Ceremonial Shield from the Solomon Islands. *Studies in Conserv.* 33 (1988) 115-125.
- [221]E. Kenndler, Neue Entwicklungen der Elektromigrationsmethoden: Hochleistungselektrophorese in Kapillaren. *Österr. Chemie-Ztg.* 12 (1988) 353.
- [222]E. Kenndler, and P. Jenner, Isotachophoresis in mixed solvents, consisting of water, methanol and dimethyl sulfoxide. Part II. Solvation Effects on the acid-base and transport properties of cation acids from ammonium ion type. *J.Chromatogr.* 390 (1987) 185-197.
- [223]E. Kenndler, and P. Jenner, Isotachophoresis in mixed solvents, consisting of water, methanol and dimethyl sulfoxide. Part I. The acid-base and migration behaviour of substituted benzene carboxylic acids. *J.Chromatogr.* 390 (1987) 169-184.
- [224]E. Kenndler, and R. Katzenberger, Isotachophoretic analysis of pharmaceutic formulations with different matrices: determination of nicotinic acid in capsules, tablets, dragees, ointments and tinctures. *Mikrochim.Acta* 1986 I (1986) 415-428.
- [225]E. Kenndler, P. Jenner, and J.F.K. Huber, Identification and quantitative estimation of organic pollutants in the exhaust air of electric ranges by two-dimensional GC and GC/MS. *Z.Anal.Chem.* 322 (1985) 198.
- [226]E. Kenndler, and E. Haidl, Mixed zone analysis in isotachophoresis with selective detection by mass spectrometry applied to the quantitation of hydrogenation products of aromatic quaternary ammonium compounds. *Z.Anal.Chem.* 322 (1985) 391.
- [227]E. Kenndler, Evaluation of isotachophoresis as a method of identification and comparison with other methods by information theory. *Analyt.Chim.Acta* 173 (1985) 239.
- [228]E. Kenndler, Applications of isotachophoresis. *Trends Anal. Chem.* 2 (1983) 202.
- [229]J.F.K. Huber, E. Matisova, and E. Kenndler, Effects of column parameters on optimization of gas chromatography/mass spectrometry. *Anal.Chem.* 54 (1982) 1297-1304.

- [230]J.F.K. Huber, E. Kenndler, W. Nyiry, and M. Oreans, Evaluation of multi-stage gas chromatography in quantitative chemical analysis. *J.Chromatogr.* 247 (1982) 211.
- [231]E. Kenndler, and D. Kaniansky, Off-line combination of isotachophoresis and mass spectrometry. *J.Chromatogr.* 209 (1981) 306.
- [232]E. Kenndler, Isotachophorese: eine Hochleistungsmethode zur Trennung von Ionen. *Österr.Chem.Zeitschr.* 3 (1981) 61.
- [233]E. Kenndler, and J.F.K. Huber, Simultaneous identification and quantitation of the food additives glutamate, guanosine 5'-monophosphate and inosine 5'-monophosphate by isotachophoresis. *Z.Lebensm.Unters.-Forsch.* 171 (1980) 292.
- [234]E. Kenndler, and J.F.K. Huber, Anwendung der analytischen Isotachophorese zur Kontrolle chemischer Produkte. *Mikrochim. Acta* 1980 (1980) 271.
- [235]E.R. Schmid, I. Fogy, F. Heresch, E. Kenndler, and J.F.K. Huber, Transfer efficiencies for the off-line combination of column liquid chromatography and mass spectrometry. *Mikrochim.Acta* 1979 (1979) 207.
- [236]J.F.K. Huber, E. Kenndler, and G. Reich, Quantitation of the information content of multi-dimensional gas chromatography and low resolution mass spectrometry in the identification of doping drugs. *J.Chromatogr.* 172 (1979) 15.
- [237]E.R. Schmid, I. Fogy, and E. Kenndler, Beitrag zur Unterscheidung von Gärungssessig, synthetischen Essig durch die Bestimmung der spezifischen 3H-Radioaktivität. *Z.Lebensm.Unters.-Forsch.* 166 (1978) 221.
- [238]E. Kenndler, and E.R. Schmid, Combination of liquid chromatography and mass spectrometry. in: J.F.K.Huber, (Ed.), *Instrumentation for high-performance liquid chromatography*, Elsevier,, Amsterdam, 1978, pp. 163-177.
- [239]J.F.K. Huber, E. Kenndler, and H. Markens, Adjustment of stationary phase selectivity in gas-liquid chromatography by using mixed liquid phases. *J.Chromatogr.* 167 (1978) 291.
- [240]E.R. Schmid, and E. Kenndler, Einfluß organischer Lösungsmittel auf die Extraktion von Thorium(IV), Uran(VI) mit substituierten sekundären und tertiären Ammoniumnitraten aus salpetersauren Lösungen. III. Aceton. *J.Radioanal.Chem.* 40 (1977) 12.
- [241]E.R. Schmid, and E. Kenndler, Einfluß organischer Lösungsmittel auf die Extraktion von Thorium(IV), Uran(VI) mit substituierten sekundären, tertiären und quartären Ammoniumnitraten aus salpetersauren Lösungen. II. Äthanol. *J.Radioanal.Chem.* 36 (1977) 317.
- [242]E.R. Schmid, I. Fogy, and E. Kenndler, Beitrag zur Unterscheidung von Gärungssessig, synthetischem Säureessig durch die Bestimmung der spezifischen 14C-Radioaktivität. *Z.Lebensm.Unters.-Forsch.* 163 (1977) 121.
- [243]E.R. Schmid, and E. Kenndler, Trennung von Thorium, Uran durch Aminextraktion aus wäßrig-methanolischen Salpetersäurelösungen. *Radiochem. Radioanal.Letters* 26 (1976) 259.
- [244]R. Schmid, E. Jünger, and E. Kenndler, Die Uran(IV)-Extraktion aus wäßrig-organischen Salpetersäurelösungen mit Trilaurylammoniumnitrat. *Radiochim.Acta* 22 (1975) 40.
- [245]E.R. Schmid, and E. Kenndler, Einfluß organischer Lösungsmittel auf die Extraktion von Thorium(IV), Uran(VI) mit substituierten sekundären, tertiären und quartären Ammoniumnitraten aus salpetersauren Lösungen. I. Methanol. *J.Radioanal.Chem.* 27 (1975) 369.

[246] E.R. Schmid, and E. Kenndler, Verteilung von Säure, Wasser, Methanol, Äthanol, Aceton zwischen gemischt wäßrig-organischen Salpetersäurelösungen und Lösungen von Trilaurylammoniumnitrat in Cyclohexan. J.Radioanal. Chem. 25 (1975) 75.

[247] E.R. Schmid, E. Jünger, and E. Kenndler, Die Extraktion von Uran (IV) aus wäßrig-organischen Salpetersäurelösungen mit dem Nitrat des sekundären Amins Amberlite LA-1. Monatsh.Chem. 104 (1973) 1560.